

## Remote Patient Monitoring (Commercial, RI Together, Direct, TMP)

Effective: November 1, 2024

<p><b>Prior Authorization Required</b> If <u>REQUIRED</u>, submit supporting clinical documentation pertinent to service request to the FAX numbers below</p>	<p>Yes <input type="checkbox"/> No <input checked="" type="checkbox"/></p>
<p><b>Notification Required</b> IF <u>REQUIRED</u>, concurrent review may apply</p>	<p>Yes <input type="checkbox"/> No <input checked="" type="checkbox"/></p>

### Applies to:

#### Commercial Products

- Harvard Pilgrim Health Care Commercial products; 800-232-0816
- Tufts Health Plan Commercial products; 617-972-9409  
CareLink<sup>SM</sup> – Refer to CareLink Procedures, Services and Items Requiring Prior Authorization

#### Public Plans Products

- Tufts Health Direct – A Massachusetts Qualified Health Plan (QHP) (a commercial product); 888-415-9055
- Tufts Health Together – MassHealth MCO Plan and Accountable Care Partnership Plans; 888-415-9055
- Tufts Health RITogether – A Rhode Island Medicaid Plan; 857-304-6404
- Tufts Health One Care – A dual-eligible product; 857-304-6304

#### Senior Products

- Harvard Pilgrim Health Care Stride Medicare Advantage; 866-874-0857
- Tufts Health Plan Senior Care Options (SCO), (a dual-eligible product); 617-673-0965
- Tufts Medicare Preferred HMO, (a Medicare Advantage product); 617-673-0965
- Tufts Medicare Preferred PPO, (a Medicare Advantage product); 617-673-0965

**Note:** While you may not be the provider responsible for obtaining prior authorization or notifying Point32Health, as a condition of payment you will need to ensure that any necessary prior authorization has been obtained and/or Point32Health has received proper notification. If notification is required, providers may additionally be required to provide updated clinical information to qualify for continued service.

## Overview

The delivery of remote patient monitoring outside of the traditional health care setting has rapidly evolved and has been proposed for patients with conditions that are at risk for decompensation and readmission. Remote patient monitoring (RPM) is the use of digital technologies to collect, measure, analyze, and/or transmit data about the physiological status of a patient outside of the traditional health setting. The data collected from these devices are then electronically transferred to providers for the purpose of monitoring disease and symptom progression remotely in order to make clinical management recommendations. RPM is considered an intervention designed to promote continuity of care and provide support for patient's and caregivers managing acute and chronic conditions; it is not intended to be an ongoing intervention for patient's whose condition(s) has stabilized.

Devices used for remote patient monitoring can include invasive and non-invasive devices, however for the purposes of this MNG, RPM refers to non-invasive devices. Examples of RPM include, but are not limited to, monitoring of blood pressure, oxygen saturation, heart rate, or weight using automated digital technology. Coverage of RPM does not apply to Continuous Glucose Monitoring (CGM) devices, Holter monitors, implantable pacemakers and defibrillators, or electroencephalograms.

Current evidence, including randomized controlled trials, meta-analyses and systematic reviews have been published pertaining to RPM and the results show a reduction in acute care use for patients with cardiovascular disease and heart

failure, however study design limitations were noted, including variability in the type of monitoring devices studied. Despite this, there is some evidence to support a reduction in the number of emergency department visits, inpatient hospitalizations, reduced mortality rates, improved patient satisfaction, and increased patient self-care management in patients with chronic obstructive pulmonary disease (COPD), and heart failure (HF). Factors contributing to readmission in patients with HF include, age, gender, body mass index (BMI), medication adherence, smoking, history of prior hospital admissions, length of hospital stay, and the presence of comorbidities such as diabetes mellitus, renal failure, coronary artery disease, and lung disease. Additionally, noncompliance was cited as a significant factor associated with HF readmission. For COPD, comorbidities, previous exacerbations and hospitalizations, and increased hospital length of stay contribute to 30- and 90-day rehospitalization rates after an index hospitalization with COPD exacerbation. Patients at the highest risk of readmission are those with multiple comorbidities, prior hospital admission, frailty, poor nutritional and low socioeconomic status.

---

## Clinical Guideline Coverage Criteria

The Plan considers remote patient monitoring (RPM), also referred to as remote physiologic monitoring, as reasonable and medically necessary when a member has a chronic or acute condition and is currently in or recently completed a skilled homecare episode of treatment or is a direct admit to the program and **ALL** of the following indications are met:

1. Member has a diagnosis of **ONE** of the following:
  - a. Chronic obstructive pulmonary disease (COPD); **or**
  - b. Heart failure (HF)
2. A physician or qualified healthcare practitioner (QHP) has documented the Member's condition is at high-risk for instability or risk of deterioration and **One** of the following:
  - a. Member has a history of two or more hospitalizations or emergency department visit(s) in the past twenty-four months for one of the conditions listed above; **or**
  - b. Member has risk factors which places the Member at risk for ED or hospitalization (e.g., recent discharge from inpatient stay or extended stay skilled nursing facility, documented poor adherence to ordered medication, or a documented history of care access challenges such as consistently missed appointments); **and**
3. Condition requires enhanced monitoring on a daily basis to assess for acute changes in clinical status and prompt intervention; **and**
4. For new Members to the physician's practice or Members not seen by the practitioner within one year, the practitioner must first conduct a face-to-face or telehealth visit with the Member to initiate RPM; **and**
5. A written order by the treating provider that specifies the medical condition and the length of time for RPM, up to 90 days
6. The Member's current/new treatment plan requires monitoring and oversight to achieve the plan of care goals; **and**
7. Remote physiologic data requirements, **All**:
  - a. The device used to collect and transmit data must meet the definition of a medical device as defined by the FDA\*; **and**
  - b. Data will be electronically collected and automatically uploaded to a secure location where the data can be available for analysis and interpretation daily by the billing practitioner; **and**
  - c. RTM data is being regularly assessed to detect acute changes in the Member's clinical status and prompt intervention.

The plan considers RPM services beyond 90 days as medically necessary when **All** of the following are met:

1. Member has complex COPD and/or HF and continues to be at high risk for hospitalization and/or ED visits; and
2. There is a physician/QHP order for the continuation of RPM; and
3. The physician/QHP is actively involved in the RPM monitoring process and is providing interpretation of the submitted Member's physiological data; and
4. The medical record contains documentation that reflects the results of the monitoring and how it is used for medical decision making.

\* The RPM device used for data collection must be a medical device, as defined by the FDA. Medical devices that digitally collect and transmit a patient's physiologic data must be reasonable and necessary for the diagnosis or treatment of the patient's illness or injury or to improve the functioning of the patient. According to the FDA, non-invasive remote monitoring devices are used to acquire patient physiological data without the need for in-clinic visits and facilitate patient management by healthcare providers while reducing the need for in-office or in-hospital services. Additionally, these devices must have the potential to be connected to a wireless network through Bluetooth, Wi-Fi, or cellular connection to transmit a patient's measurements directly to their healthcare provider or other monitoring entities. The device used must provide secure, HIPAA-compliant transmission of the data.

## Limitations

The Plan considers remote patient monitoring as not medically necessary under the following conditions:

1. Member and/or caregiver is not capable of using the equipment
2. The RPM device itself (including any additional apps, software, digital interfaces, etc.) is not covered
3. Use of smartwatches and fitness trackers

## Codes

The following code(s) do not require prior authorization: (Note: Monitoring must occur over at least 16 days of a 30-day period. These codes should not be reported for monitoring if the duration is less than 16 days in order for CPT codes 99453 and 99454 to be billed)

**Table 1: CPT/HCPCS Codes**

Code	Description
99091	Collection and interpretation of physiologic data (eg, ECG, blood pressure, glucose monitoring) digitally stored and/or transmitted by the patient and/or caregiver to the physician or other qualified health care professional, qualified by education, training, licensure/regulation (when applicable) requiring a minimum of 30 minutes of time, each 30 days
99453	Remote monitoring of physiologic parameter(s) (eg, weight, blood pressure, pulse oximetry, respiratory flow rate), initial; set-up and patient education on use of equipment
99454	Remote monitoring of physiologic parameter(s) (eg, weight, blood pressure, pulse oximetry, respiratory flow rate), initial; device(s) supply with daily recording(s) or programmed alert(s) transmission, each 30 days
99457	Remote physiologic monitoring treatment management services, clinical staff/physician/other qualified health care professional time in a calendar month requiring interactive communication with the patient/caregiver during the month; first 20 minutes
99458	Remote physiologic monitoring treatment management services, clinical staff/physician/other qualified health care professional time in a calendar month requiring interactive communication with the patient/caregiver during the month; each additional 20 minutes (List separately in addition to code for primary procedure)

## List of Medically Necessary ICD-10 Codes

### References:

1. American College of Physicians (ACP) Remote patient monitoring billing, coding and regulations information. Available at URL address: [Remote Patient Monitoring | ACP \(acponline.org\)](https://www.acponline.org/remote-patient-monitoring). Accessed October 16, 2024.
2. American Heart Association. AHA Guidance: Using Remote Patient Monitoring Technologies for Better Cardiovascular Disease Outcomes. 2019. <https://www.heart.org/-/media/files/about-us/policy-research/policy-positions/clinical-care/remote-patient-monitoring-guidance-2019.pdf?la=en&hash=A98793D5A043AB9940424B8FB91D2E8D5A5B6BEB>. Accessed October 16, 2024.
3. Burke RE, Schnipper JL, Williams MV, et al. The HOSPITAL Score Predicts Potentially Preventable 30-Day Readmissions in Conditions Targeted by the Hospital Readmissions Reduction Program. *Med Care*. 2017;55(3):285-290. doi:10.1097/MLR.0000000000000665.
4. Center for Connected Health Policy Remote Patient Monitoring. <https://www.cchpca.org/about/about-telehealth/remote-patient-monitoring-rpm>. Accessed October 16, 2024.
5. Centers for Medicare & Medicaid Services (CMS) Medicare Learning Network (MLN) Telehealth Services Fact Sheet June 2023 MLN901705 - Telehealth Services (cms.gov). Accessed October 16, 2024.
6. Commonwealth of Massachusetts Executive Office of Health and Human Services Office of Medicaid. Transmittal Letter CHC-122. [\\*Transmittal Letter \(mass.gov\)](https://www.mass.gov/info-details/transmittal-letter-chc-122). July 24, 2024. Accessed August 8, 2024.
7. Commonwealth of Massachusetts Executive Office of Health and Human Services Administrative Bulletin 22-09. [The Commonwealth of Massachusetts](https://www.mass.gov/info-details/transmittal-letter-chc-122). January 1, 2022. Accessed August 7, 2024
8. Coutu FA, Iorio OC, Ross BA. Remote patient monitoring strategies and wearable technology in chronic obstructive pulmonary disease. *Front Med (Lausanne)*. 2023;10:1236598. Published 2023 Aug 17. doi:10.3389/fmed.2023.1236598.
9. Federal Register, A Rule by the Centers for Medicare & Medicaid Services on <https://www.govinfo.gov/content/pkg/FR-2020-12-28/pdf/2020-26815.pdf>. Accessed October 16, 2024.
10. Global Initiative for Chronic Obstructive Lung Disease. Global strategy for prevention, diagnosis and management of COPD: 2024 report. Global Initiative for Chronic Obstructive Lung Disease; 2024. Available from: [2024 GOLD](https://goldcopd.org/)

[Report - Global Initiative for Chronic Obstructive Lung Disease - GOLD \(goldcopd.org\)](#). Accessed October 16, 2024.

11. Jayaram NM, Khariton Y, Krumholz HM, et al. Impact of Telemonitoring on Health Status. *Circ Cardiovasc Qual Outcomes*. 2017;10(12):e004148. doi:10.1161/CIRCOUTCOMES.117.004148.
12. Kew KM, Cates CJ. Home telemonitoring and remote feedback between clinic visits for asthma (Review). *Cochrane Database of Systematic Reviews* 2016, Issue 8. Art. No:CD011714.
13. Khanijahani A, Akinci N, Quitiquit E. A Systematic Review of the Role of Telemedicine in Blood Pressure Control: Focus on Patient Engagement. *Curr Hypertens Rep*. 2022;24(7):247-258. doi:10.1007/s11906-022-01186-5.
14. Kitsiou S, et al. Effects of home telemonitoring interventions on patients with chronic heart failure: an overview of systematic reviews. *J Med Internet Res* 2015, Vol 17, Iss 3, e63.
15. Kruse C, et al. Telemonitoring to manage chronic obstructive pulmonary disease: systematic literature review. *JMIR Medical Informatics*. Vol 7, No 1 (2019):Jan-Mar.
16. Lee PA, Greenfield G, Pappas Y. The impact of telehealth remote patient monitoring on glycemic control in type 2 diabetes: a systematic review and meta-analysis of systematic reviews of randomised controlled trials. *BMC Health Serv Res*. 2018;18(1):495. Published 2018 Jun 26. doi:10.1186/s12913-018-3274-8.
17. Lu JK, Sijm M, Janssens GE, Goh J, Maier AB. Remote monitoring technologies for measuring cardiovascular functions in community-dwelling adults: a systematic review [published online ahead of print, 2023 May 18]. *Geroscience*. 2023;1-12. doi:10.1007/s11357-023-00815-4.
18. Mayo Clinic. Home Telemonitoring in Patients at High Risk for Readmission. NCT02136186. Last updated November 10, 2020. Available at: <https://clinicaltrials.gov/ct2/show/NCT02136186>. Accessed on October 16, 2024.
19. Mehr DR, Wakefield D, Caligiuri F, et al. Using Health Information Technology (HIT) to Improve Ambulatory Chronic Disease Care. Agency for Healthcare Research and Quality (AHRQ) Grant ID: HS017035. Inclusive Dates: 9/13-2007- 8/31/11. Available at: <https://digital.ahrq.gov/sites/default/files/docs/publication/r18hs017035-mehr-final-report-2011.pdf>. Accessed on October 16, 2024.
20. Milani RV, et al. Improving hypertension control and patient engagement using digital tools. *Am J Med*. 2017 Jan;130(1):14-20.
21. Nick JM, Roberts LR, Petersen AB. Effectiveness of telemonitoring on self-care behaviors among community-dwelling adults with heart failure: a quantitative systematic review. *JBIE Evid Synth*. 2021;19(10):2659-2694. doi:10.11124/JBIES-20-00329.
22. Scholte NTB, Gürgöze MT, Aydin D, et al. Telemonitoring for heart failure: a meta-analysis. *Eur Heart J*. 2023;44(31):2911-2926. doi:10.1093/eurheartj/ehad280.
23. Singh G, Zhang W, Kuo YF, Sharma G. Association of Psychological Disorders With 30-Day Readmission Rates in Patients With COPD. *Chest*. 2016;149(4):905-915. doi:10.1378/chest.15-0449.
24. Taylor ML, Thomas EE, Snoswell CL, Smith AC, Caffery LJ. Does remote patient monitoring reduce acute care use? A systematic review. *BMJ Open*. 2021;11(3):e040232. Published 2021 Mar 2. doi:10.1136/bmjopen-2020-040232.
25. Totten AM, Womack DM, Eden KB, McDonagh MS, Griffin JC, Grusing S, Hersh WR. Telehealth: Mapping the Evidence for Patient Outcomes From Systematic Reviews [Internet]. Rockville (MD): Agency for Healthcare Research and Quality (US); 2016 Jun. Report No.: 16-EHC034-EF. PMID: 27536752..
26. U.S. Food and Drug Administration, U.S. Department of Health and Human Services Food and Drug Administration [Enforcement Policy for Non-Invasive Remote Monitoring Devices Used to Support Patient Monitoring \(fda.gov\)](#) October 19, 2023. Accessed October 24, 2024.
27. U.S. Government Accountability Office. Health care: Telehealth and remote patient monitoring use in the Medicare and selected federal programs. <https://www.gao.gov/assets/gao-17-365.pdf>. Accessed January 4, 2024.
28. Veenis JF, Radhoe SP, Hooijmans P, Brughts JJ. Remote Monitoring in Chronic Heart Failure Patients: Is Non-Invasive Remote Monitoring the Way to Go?. *Sensors (Basel)*. 2021;21(3):887. Published 2021 Jan 28. doi:10.3390/s21030887.

---

## Approval And Revision History

January 17, 2024: Reviewed by the Medical Policy Approval Committee (MPAC) as a new coverage guideline for Remote Patient Monitoring for Members with COPD and Heart Failure effective May 1, 2024

Subsequent endorsement date(s) and changes made:

- September 19, 2024: Reviewed by the Medical Policy Approval Committee (MPAC) with an effective date of November 1, 2024: Renamed to Remote Monitoring for Commercial, RI Together, Direct, TMP products; minor content changes made for clarity and added new criteria for remote patient monitoring beyond 90 days and for new members to the physician's practice or in the event that a member has not seen the practitioner within one year, the practitioner must first conduct a face-to-face or telehealth visit with the member to initiate remote patient monitoring

---

## Background, Product and Disclaimer Information

Medical Necessity Guidelines are developed to determine coverage for benefits and are published to provide a better understanding of the basis upon which coverage decisions are made. We make coverage decisions using these guidelines, along with the Member's benefit document, and in coordination with the Member's physician(s) on a case-by-case basis considering the individual Member's health care needs.

Medical Necessity Guidelines are developed for selected therapeutic or diagnostic services found to be safe and proven effective in a limited, defined population of patients or clinical circumstances. They include concise clinical coverage criteria based on current literature review, consultation with practicing physicians in our service area who are medical experts in the particular field, FDA and other government agency policies, and standards adopted by national accreditation organizations. We revise and update Medical Necessity Guidelines annually, or more frequently if new evidence becomes available that suggests needed revisions.

For self-insured plans, coverage may vary depending on the terms of the benefit document. If a discrepancy exists between a Medical Necessity Guideline and a self-insured Member's benefit document, the provisions of the benefit document will govern. For Tufts Health Together (Medicaid), coverage may be available beyond these guidelines for pediatric members under age 21 under the Early and Periodic Screening, Diagnostic and Treatment (EPSDT) benefits of the plan in accordance with 130 CMR 450.140 and 130 CMR 447.000, and with prior authorization.

Treating providers are solely responsible for the medical advice and treatment of Members. The use of this guideline is not a guarantee of payment or a final prediction of how specific claim(s) will be adjudicated. Claims payment is subject to eligibility and benefits on the date of service, coordination of benefits, referral/authorization, utilization management guidelines when applicable, and adherence to plan policies, plan procedures, and claims editing logic.